

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**

STIC-ILL

V. No 3/19

From: Holleran, Anne  
Sent: Wednesday, March 19, 2003 2:27 PM  
To: STIC-ILL  
Subject: refs. for 09/302,434

437248

Please send copies of the following papers:

1. Gussow Methods in Enzymology (1991) 203: 99-121
2. Bosslet Eur. J. Nucl. Med. (1988) 14: 523-528

Anne Holleran  
AU: 1642  
Tel: 308-8892  
RM: 8e03

mailbox: 8e12

6/10/03

15810177

## Quantitative considerations supporting the irrelevance of circulating serum CEA for the immunoscintigraphic visualization of CEA expressing carcinomas

K. Bosslet<sup>1</sup>, A. Steinsträsser<sup>2</sup>, A. Schwarz<sup>2</sup>, H.P. Harthus<sup>1</sup>, G. Lüben<sup>1</sup>, L. Kuhlmann<sup>2</sup>, and H.H. Sedlacek<sup>1</sup>

<sup>1</sup> Research Laboratories of Behringwerke AG, P.O. Box 1140, D-3550 Marburg/Lahn, Federal Republic of Germany

<sup>2</sup> Radiochemical Laboratory of Hoechst AG, Strooßstrasse, D-6230 Frankfurt/Main 80, Federal Republic of Germany

**Abstract.** Starting from the phenomenon that the amount of circulating CEA in patients' sera did not significantly influence immunoscintigraphic visualization of CEA expressing tumors, we built up an in vitro model to explain this phenomenon. Blocking experiments in this model system showed that the CEA specific MAbs BW 431/26 and BW 431/31 could not be inhibited in their binding to cell associated CEA, if they were preincubated with a 20 molar excess of serum CEA. In contrast, the CEA-NCA cross reactive MAbs could be inhibited in their binding to tumor associated CEA under identical conditions. These data combined with western blotting analysis of patients' sera and affinity constant determinations argue that conformational changes in serum CEA cause a decreased affinity of the CEA specific MAbs to serum CEA allowing a preferential binding to tumor associated CEA.

**Key words:** CEA - Conformational change - MAb - Immunoscintigraphy

Since the first description of carcinoembryonic antigen (CEA) (Gold and Freedman 1965a, b) until today, where the primary structure of CEA deduced from the cDNA sequence (Oikawa et al. 1987a) is known, CEA became important as a tumor marker for the treatment control of colorectal carcinomas (Egan et al. 1977; Fuks et al. 1980) as well as for the immunoscintigraphic visualization of CEA expressing carcinomas (Mach et al. 1981). In particular, the relatively new method of immunoscintigraphy demands monoclonal antibodies (MAb) binding to epitopes on CEA not detectable on the various nonspecific cross-reacting antigens (NCAs) expressed on normal tissues (von Kleist et al. 1972; Rogers 1983). A so called anti CEA antibody having

strong cross-reactivity to human granulocytes and erythrocytes, i.e. NCA, was not able to localize human colorectal carcinomas in vivo, but induced fever, rigors, emesis and a short term granulocytopenia (Dillman et al. 1984). Also, those MAbs which are selective for protein epitopes on CEA (Buehgeger et al. 1984; Bosslet et al. 1985) but not cross-reactive with epitopes on NCAs or Y or X type carbohydrate moieties on CEA (Nichols et al. 1985) are, after i.v. injection into the patient, primarily confronted with high amounts of circulating CEA. The following in vitro experiments try to explain why successful immunoscintigraphic localization of colorectal carcinomas using MAbs binding to CEA are possible with a sensitivity of up to 90% as shown in a multicenter study (100 patients) in the FRG using the MAb BW 431/31, coupled via DTPA to <sup>111</sup>In (Bosslet et al. 1987).

### Materials and methods

**Tumor cell lines.** The LoVo adenocarcinoma cell line of the colon was obtained from the American Type Culture Collection (ATCC, 12301 Parklawn Drive, Rockville, MD 20852, USA/ATCC-No.: CCL 229). It was routinely passaged in RPMI-1640 with 10% FBS. Confluent cultures were treated for 3 min with 0.25% trypsin 0.02 M EDTA and subcultured at a ratio of 1:5

**Monoclonal antibodies.** The MAbs BW 374/14, BW 250/183, BW 431/26 and BW 431/31, all of IgG<sub>1</sub>-isotype, recognize different epitopes on CEA (Bosslet et al. 1985, 1987). Briefly, the MAbs BW 431/26 and BW 431/31 are selective for CEA, whereas the MAbs BW 374/14 or BW 250/183 bind to NCA55 (=non specific cross-reacting antigen), NCA95 and CEA or NCA95 and CEA, respectively. MAb BW 494 binds to a pancreatic, gastric and colonic carcinoma associated mucin molecule different from CEA (Bosslet et al. 1986). MAb BW 575 is selective for small cell lung carcinomas and does not show any significant binding to gastrointestinal tract carcinomas (Bosslet 1987).

**Preparation of peripheral blood granulocytes.** Human peripheral blood granulocytes were isolated from heparin blood by treating the Ficoll-Hypaque pellet (Boyum 1976) for 5 min at 37° C with 0.87% NH<sub>4</sub>Cl to lyse erythrocytes.

Offprint requests to: K. Bosslet

### Abbreviations used

CEA carcinoembryonic antigen; NCA nonspecific cross-reacting antigen; DTPA diethylene triamine pentaacetic acid; FBS fetal bovine serum; EDTA ethylene diamine tetraacetic acid; PCA perchloric acid; R.T. room temperature; PBS phosphate buffered saline; BSA bovine serum albumin; SDS-PAGE sodium dodecyl-sulfate polyacrylamide gel electrophoresis

**Preparation of granulocyte extracts.** The erythrocyte depleted Ficoll-Hypaque pellet was solubilized by adding 0.5% Na-deoxycholate followed by 12 h head over head rotation at 4° C. After ultracentrifugation at 30000 g for 30 min at 4° C, aliquots of the supernatant corresponding to  $2 \times 10^5$  granulocyte equivalents were transferred to individual lanes for SDS-PAGE and western blotting.

**Radiolabelling of MABs.** Purified MABs (Ey et al. 1978) were radiolabelled with  $^{125}\text{I}$  using the iodogen method (Fraker and Speck 1978; Markwell and Fox 1978). MAB proteins were diluted in PBS containing 0.1% BSA (PBS-BSA).

The specific activities obtained for the different MABs were:  
 BW 374/14: 3.9  $\mu\text{Ci}/\mu\text{g}$ , BW 250/183: 3.5  $\mu\text{Ci}/\mu\text{g}$   
 BW 431/31: 3.9  $\mu\text{Ci}/\mu\text{g}$ , BW 431/26: 2.9  $\mu\text{Ci}/\mu\text{g}$   
 BW 494: 4.03  $\mu\text{Ci}/\mu\text{g}$ , BW 575: 2.59  $\mu\text{Ci}/\mu\text{g}$

**Blocking assay.**  $^{125}\text{I}$  labelled MAB protein (100  $\mu\text{l}$ ) diluted in PBS-BSA was mixed with CEA containing serum (100  $\mu\text{l}$ ) or purified CEA isolated from CEA expressing human colon carcinomas and incubated for 90 min at 37° C in round bottom 96 well microtiter plates (Nunc: 1-63320). Thereafter,  $10^6$  LoVo cells in 25  $\mu\text{l}$  PBS-BSA which were detached from plastic by incubation with 0.02% EDTA for 5 min, were added to the MAB serum or MAB-CEA mixture. After 1 h incubation at 37° C, the microtiter plates were spun and the pellets were washed 3 times with PBS. The washed cell pellet was resuspended in 125  $\mu\text{l}$  and 100  $\mu\text{l}$  thereof and counted in a well type  $\gamma$ -counter to determine the cell bound radioactivity.

**Western blotting analysis.** The methodology was performed essentially as described by Towbin and Gordon (1979), but instead of radiolabelled second antibody, the ABC vector kit (Vector Burlingame) was used.

**Scatchard analysis.** The affinity of the MABs to tumor cells was evaluated according to Walker (1977), based on the original papers of Scatchard (1949) and Berson and Yalow (1959). As antigenic material collagenized and formaldehyde fixed cells derived from nude mouse xenografts transplanted with the CEA expressing CoCa<sup>2</sup> colonic carcinoma were used.

A constant amount of these cells is added to each vial of a dilution series of the respective MAB. This mixture is incubated for 4 h at room temperature, the cells were spun down, washed and the cell bound activity determined.

The bound to free ratio is plotted against concentration of cell bound antibody molecules using the following corrections (Steinsträsser et al. 1987): (1) The unspecific bound part (U) of antibodies (determined with irrelevant cells or by an iteration procedure during the evaluation) is subtracted. (2) The free part of antibodies (F) consists only of immune reactive molecules using the formula  $F = r \times T - (B - U)$ , where T is the total activity and r the immune reactive fraction of the preparation, determined separately according to Steinsträsser and Schwarz (1986).

The inverse of the slope of the described plot gives the affinity constant (l/mol).

**CEA assay.** The amount of CEA present in patients' sera was evaluated using the commercially available Behring Enzygnost CEA assay.

## Results

### Binding inhibition experiments on LoVo cells

Incubation of  $10^6$  CEA expressing living LoVo colon carcinoma cells with 2–3 ng (10000 cpm) of the various MAB proteins for 60 min at 37° C results in the binding of 0.2–0.3 ng (1000 cpm) to the membrane of the cells. Preincubation of the various MABs ( $\approx 10000$  cpm) for 90 min at 37° C with colon carcinoma patients' sera (Fig. 1) containing high levels (a–d), medium levels (e–f), or low levels (g–j) of CEA had the following effects on MAB binding to the cells (Fig. 1):

The binding of MABs BW 431/26 and BW 431/31 to the LoVo cells is only weakly inhibited despite a 20 fold molar excess of soluble CEA (a–d). In contrast, the binding of MABs BW 250/183 and BW 374/14 to the LoVo cells is efficiently inhibited under identical conditions (a–d). Marginal binding inhibition was observed with all anti CEA MABs with medium (e–f) or low (g–j) CEA containing sera.

Purified CEA (Krupey et al. 1968) isolated from a colon carcinoma primary tumor did not inhibit the binding of MABs BW 431/26 and BW 431/31 to the LoVo cells as efficiently as it inhibited the binding of MABs BW 374/14 and BW 250/183 (k).

The positive control MAB BW 494 binding to a high molecular weight mucin epitope (Bosslet et al. 1986) not detectable on CEA could not be inhibited in its binding to the LoVo cells, neither by CEA containing sera (a–j) nor by purified CEA (k). The binding of the anti small cell lung carcinoma MAB BW 575 (Bosslet et al. 1987) was in the range 1%–3% of the binding obtained with the anti CEA MABs or MAB BW 494 (data not shown). Purified human serum albumin (l) as well as phosphate buffered saline (m) showed no inhibition of binding with any MAB.

In summary, the binding of the anti CEA-NCA MABs BW 374/14 and BW 250/183 is equally effectively inhibited by CEA containing sera and purified CEA, whereas the binding of the anti CEA MABs BW 431/26 and BW 431/31 is neither inhibited by the purified CEA isolated from a primary colonic carcinoma nor by the CEA containing sera.

### Western blotting analysis of CEA containing patient's sera

To investigate whether the reduced inhibition mediated by the CEA containing sera on the binding of the CEA specific MABs BW 431/26 and BW 431/31 to the LoVo colonic carcinoma cell line is due to the absence of intact CEA in the sera, we performed a western blotting analysis after perchloric acid (PCA) extraction with the same sera and CEA preparation as used in the preceding inhibition experiments. The data presented in Fig. 2A, B, C, D show that all four MABs bound to purified CEA attached to nitrocellulose filter (1  $\mu\text{g}$ ) to a similar extent, visualizing a 180 kDa band (lane k, Fig. 2, A, B, C, D). The CEA specific MABs BW 431/26 and BW 431/31 did not bind to PCA extracts from colonic carcinoma patients' sera (Fig. 2A, B, lane a, b, c, d), whereas the CEA-NCA cross-reactive MABs visualize the 180 kDa CEA band (Fig. 2, C, D lane a, b, c, d). Neither MAB bound to PCA extracts of a control serum (Fig. 2A, B, C, D lane g). As a positive control for the binding potential of MABs BW 374/14 and BW 250/183 to NCA95, they were reacted with Na-deoxycholate extracts from human peripheral blood granulocytes. In lane o, Fig. 2C, D, the NCA band was significantly stained, where-




Fig. 1a-m. Inhibition of CEA after blocking with CEA (k), HSA (l), and PBS (m).

as the 2 CEA bands did not show inhibition (Fig. 2A, B).

In summary, patients' sera detected by the CEA-NCA MABs BW 374/14 and BW 250/183 contained CEA from a similar source to a similar extent, irrelevant MAB binding to the cells (data not shown).

### Estimation of membrane bound CEA

To exclude various MABs served in the experiments, the anti CEA MABs BW 374/14 and BW 250/183 were used. Methods. A fixed cells were used. Table 1.

olon carc-  
ous MAb  
inding of  
ills. Prein-  
or 90 min  
g. 1) con-  
low levels  
b binding

431/31 to  
a 20 fold  
ie binding  
oVo cells  
ns (a-d).  
anti CEA  
ning sera.  
m a colon  
inding of  
ills as effi-  
74/14 and

to a high  
1986) not  
s binding  
sera (a-j)  
nti small  
1987) was  
h the anti  
. Purified  
buffered  
ny MAb.  
A MAb  
inhibited  
ereas the  
W 431/31  
d from a  
ning sera.

t's sera

diated by  
A specific  
lonic car-  
CEA in  
ysis after  
sera and  
n experi-  
how that  
nitrocel-  
180 kDa  
ic MAb  
extracts  
3, lane a,  
as visual-  
b, c, d).  
ol serum  
l for the  
50/183 to  
extracts  
1 lane o,  
d, where-




Fig. 1 a-m. In the binding inhibition experiment black columns represent the percentage of the respective MABs bound to cell associated CEA after blocking with sera from patients with high (a-d), medium (e, f) or low (g-j) CEA content in comparison to purified swimming CEA (k), HSA (l) or PBS (m)

as the 2 CEA specific MABs BW 431/26 and BW 431/31 did not show any binding to granulocyte NCA (lane o, Fig. 2 A, B).

In summary, these data indicate that colonic carcinoma patients' sera contained intact 180 kDa CEA which was detected by the CEA-NCA cross-reactive MABs, but not by the CEA specific MABs. No NCA could be detected in patients' sera by neither MAB. But granulocyte associated NCA95 (Buchegger et al. 1984) was detected by MABs BW 374/14 and BW 250/183. In contrast purified CEA from a colonic carcinoma primary tumor was detected to a similar extent by the four specific MABs used. The irrelevant MABs BW 494 and BW 575 did not show any binding to purified CEA or CEA containing patients' sera (data not shown).

#### Estimation of the *in vitro* binding affinity of MABs to cell membrane bound CEA

To exclude that simple differences in the affinity of the various MABs to CEA were responsible for the results observed in the binding inhibition and western blotting experiments, the affinity constants of the MABs to tissue associated CEA were determined as detailed in Material and Methods. As antigen source, collagenized, formaldehyde fixed cells from colon carcinoma nude mouse xenografts were used. The affinity constants calculated are shown in Table 1.

Original scatchard plots from a representative experiment are shown in Fig. 3 for MAB BW 431/26 (A), BW 431/31 (B) and BW 250/183 (C). In Table 1 affinity constants evaluated from various experiments are given, supporting the medium affinity constant values mentioned above.

#### Discussion

The paper deals with the so far unexplained phenomenon that CEA specific MABs are able to localize CEA expressing tumors in the patient despite significant amounts of circulating CEA in the blood. In the binding inhibition experiments we could show that MABs directed to CEA specific epitopes on CEA (MAB BW 431/26 and BW 431/31) are only marginally blocked by CEA containing sera. The CEA content of these sera was in 20fold molar excess (500 ng CEA  $\leftrightarrow$  2 ng MAB) compared to the radiolabelled MABs used in the binding assays.

In contrast, the CEA-NCA cross-reactive MABs could be more efficiently blocked under identical conditions. These findings could be explained by the presence of NCA in the CEA containing sera mediating the more efficient blocking of the CEA-NCA cross-reactive MABs, or by higher affinities of the CEA-NCA MABs to circulating CEA. The western blotting analysis performed thereafter showed that the CEA containing sera did not contain detectable amounts of NCA, arguing against the first explanation. In contrast, NCA from normal human granulocytes




Fig. 2. Photograph of 4 blots independently blotted by means of 10% SDS-PAGES immunostained with A MAb BW 431/26, B MAb BW 431/31, C MAb BW 250/183, D MAb BW 374/14. Lanes a-d represent PCA extracts from patients sera containing CEA (compare Fig. 1), lane g represents control serum, lane k represents purified CEA, lane o represents Na-deoxycholate extracts from normal human granulocytes ( $2 \times 10^5$  granulocyte equivalents/lane). Lane z visualizes the rainbow protein molecular weight markers (Amersham)

Table 1. Determination of affinity-constant by scatchard evaluation


MAb	Number of determinations	$K_a \pm sd$	range
431/31	17	$3 \pm 2 \times 10^8$	$1 \times 10^8 \dots 8 \times 10^8$
250/183	6	$2 \pm 1 \times 10^9$	$1 \times 10^9 \dots 4 \times 10^9$
431/26	5	$9 \pm 12 \times 10^9$	$1 \times 10^9 \dots 3 \times 10^{10}$

could be detected under the same experimental conditions with MAbs BW 250/183 and BW 374/14. Furthermore, the western blot data indicated that the CEA-specific MAbs (BW 431/26 and BW 431/31) as well as the CEA-NCA cross-reactive MAbs (BW 250/183 and BW 374/14) were able to detect CEA purified from primary tumors attached to nitrocellulose filter to a similar extent.

Therefore, the second explanation that the CEA-NCA cross-reactive MAbs have a higher affinity to circulating CEA than the CEA-specific ones is probably correct. The scatchard plot data showed that the CEA-specific MAb

BW 431/26 had the highest affinity (in the range  $10^{10}$  l/mol) if cell bound CEA is used as epitope bearing antigen. The reduced binding, especially of this MAb to swimming CEA as evidenced by the binding inhibition experiments could be explained by differences in the conformation between cell bound and circulating CEA. Especially the class five epitopes (Kuroki et al. 1987) which are unique to CEA are known to be weakly immunogenic if purified CEA is used as immunogen. Furthermore, the affinity of those MAbs to CEA is mostly in the range of  $10^8$  l/mol, but MAb BW 431/26, having an affinity constant of  $9 \times 10^9$  l/mol, is an exception allowing successful immunoscintigrams of colorectal carcinomas 4–6 h after i.v. application of the  $^{99m}Tc$ -MAb-immunoconjugate in the patient (A. Schwarz manuscript in preparation). The success of the immunoscintigraphic evaluation is independent of the CEA content in the patients' serum as revealed in a multicenter study performed with MAb BW 431/31 (Bosslet et al. 1987).

This clinical finding is mimicked by our in vitro binding inhibition model. We confronted our radiolabelled MAbs for 90 min with up to 500 ng ( $1.6 \times 10^{12}$  molecules) CEA from CEA containing sera. This is a 20fold molar excess compared to the 2 ng of MAb ( $8 \times 10^{10}$  molecules). There-








Fig. 3A-C. Scatchard plots are depicted where the bound (B) to free ratio (F) was plotted against the cell bound MAb concentration. A-C represent plots obtained for MAb BW 431/26, BW 431/31 and BW 250/183 respectively

ots  
means of  
ostained  
B MAb  
50/183, D  
-d  
m patients  
ipare  
ontrol  
urified  
1-  
1 normal  
0<sup>5</sup>  
ne). Lane z  
tein

10<sup>10</sup> l/mol)  
gen. The  
ing CEA  
its could  
between  
class five  
CEA are  
A is used  
e MAb  
ut MAb  
0<sup>9</sup> l/mol,  
grams of  
n of the  
Schwarz  
unoscint  
-tent in  
udy per-

binding  
d MAb  
s) CEA  
r excess  
There-

after, to this mixture  $10^6$  LoVo cells were added bearing between  $10^{10}$ – $10^{11}$  CEA epitopes ( $10^4$ – $10^5$  epitopes/cell) and incubated for another h. The binding data showed that especially the CEA specific MABs BW 431/26 and BW 431/31 bound very efficiently to the CEA expressing cells arguing for a preferential binding of these MABs to cell bound CEA. For scintigraphic purposes with anti CEA MABs, patients receive 2 mg of purified MAB resulting in a concentration of 670 ng MAB/ml ( $2.7 \times 10^{12}$  molecules/ml). The serum CEA concentrations in colonic carcinoma patients are mostly in the range 1–3 ng CEA/ml (maximally  $1 \times 10^{13}$  molecules/ml) yielding maximally a 4fold molar excess of CEA compared to the MAB in the serum. This molar excess of circulating CEA does not seem to be high enough to block the binding of the MAB to the tumor associated CEA epitopes. Despite our detailed knowledge about CEA, i.e. the cDNA sequence (Oikawa et al. 1987a), the immunoglobulin domain like structure (Oikawa et al. 1987b), the N linked carbohydrate chains (Chandrasekaran et al. 1983) and the  $\text{NH}_2$  terminal amino acid sequence (Engvall et al. 1978), the exact structure of MAB defined epitopes as well as the MAB detected conformational differences between serum CEA and tumor associated CEA remain to be elucidated.

**Acknowledgement.** The technical assistance of H. Schmidt and N. Döring as well as the secretarial assistance of I. Gimbel and S. Lehnert are greatly appreciated.

## References

- Berson SA, Yalow RS (1959) Quantitative aspects of the reaction between insulin-binding antibody. *J Clin Invest* 38:1996–2016
- Bosslet K (1987) Monoclonal antibodies suited for the immunoscintigraphic differential diagnosis of human lung cancer. Meeting Abstract: 3rd International Congress on Hormones and Cancer, September 1987, Hamburg, FRG
- Bosslet K, Lüben G, Schwarz A, Hundt E, Harthus HP, Seiler FR, Muhrer C, Klöppel G, Kayser K, Sedlacek HH (1985) Immunohistochemical localization and molecular characteristics of three monoclonal antibody-defined epitopes on carcinoembryonic antigen (CEA). *Int J Cancer* 36:75–84
- Bosslet K, Kern HF, Kanzy EJ, Steinstraesser A, Schwarz A, Lüben G, Schorlemmer HU, Sedlacek HH (1986) A monoclonal antibody with binding and inhibiting activity towards human pancreatic carcinoma cells. *Cancer Immunol Immunother* 23:185–191
- Bosslet K, Schwarz A, Steinstraesser A, Kuhlmann L, Seidel L, Schulz G (1988) Immunsintigraphie kolorektaler Karzinome und spezifische Immuntherapie von Pankreaskarzinomen mittels monoklonaler Antikörper. *Wiener Med Wschr* 11:255–257
- Boyum A (1976) Isolation of lymphocytes, granulocytes and macrophages. *Scand J Immunol* 5:9–15
- Buchegger F, Schreyer M, Carrel S, Mach JP (1984) Monoclonal antibodies identify a CEA cross-reacting antigen of 95 kD (NCA-95) distinct in antigenicity and tissue distribution from the previously described NCA of 55 kD. *Int J Cancer* 33:643–649
- Chandrasekaran EV, Davila M, Nixon DW, Goldfarb M, Mendicino J (1983) Isolation and structures of the oligosaccharide units of carcinoembryonic antigen. *J Biol Chem* 258:7213–7222
- Dillman RO, Beauregard JC, Sobol RE, Royston I, Bartholomew RM, Hagan PS, Halpern SE (1984) Lack of radioimmunodetection and complications associated with monoclonal anticarcinoembryonic antigen antibody. *Cancer Res* 44:2213–2218
- Egan ML, Engvall E, Ruoslahti EI, Todd CW (1977) Detection of circulating tumor antigens. *Cancer Res* 40:458–466
- Engvall E, Shively JE, Wrann M (1978) Isolation and characterization of the normal cross-reacting antigen: Homology of its  $\text{NH}_2$ -terminal amino acid sequence with that of carcinoembryonic antigen. *Proc Nat Acad Sci USA* 75:1670–1674
- Ey PL, Prowse SJ, Jenkin CR (1978) Isolation of pure  $\text{IgG}_1$ ,  $\text{IgG}_{2a}$ ,  $\text{IgG}_{2b}$  immunoglobulins from mouse serum using protein A-sepharose. *Immunochimistry* 15:429–436
- Fraker PJ, Speck JC (1978) Protein and cell membrane iodinations with a sparingly soluble chloroamide, 1,3,4,6-tetra chloro-3,6-diphenyl glycoluril. *Biochem Biophys Res Comm* 80:849–857
- Fuks A, Shuster J, Gold P (1980) Theoretical and practical considerations of the utility of the radioimmunoassay for carcinoembryonic antigen in clinical medicine. In: Sell S (ed) *Cancer markers, Human*, Clifton NJ, pp 315–327
- Gold P, Freedman SO (1965a) Demonstration of tumor-specific antigens in human colonic carcinomata by immunological tolerance and absorption techniques. *J Exp Med* 121:439–462
- Gold P, Freedman SO (1965b) Specific carcinoembryonic antigens of the human digestive system. *J Exp Med* 122:467–480
- Kleist S von, Chavanel G, Burtin P (1972) Identification of an antigen from normal human tissue that cross-reacts with the carcino-embryonic antigen. *Proc Nat Acad Sci (Wash)* 69:2492–2494
- Krupey J, Gold P, Freedman SO (1968) Physicochemical studies of the carcinoembryonic antigen of the human digestive system. *J Exp Med* 128:387–398
- Kuroki M, Arakawa F, Higachi H, Matsunaga A, Okamoto N, Takakura K, Matsuoka Y (1987) Epitope mapping of the carcinoembryonic antigen by monoclonal antibodies and establishment of a new improved radioimmunoassay system. *Jpn J Cancer Res* 78:386–396
- Mach JP, Buchegger F, Forni M, Ritschard J, Berche C, Lumbroso J-D, Schreyer M, Girardet C, Accolla RC, Carrel S (1981) Use of radiolabelled monoclonal anti-CEA antibodies for the detection of human carcinomas by external photoscanning and tomoscintigraphy. *Immunology today* 2:239–249
- Markwell MAK, Fox CF (1978) Surface-specific iodination of membrane proteins of viruses and eucaryotic cells using 1,3,4,6-tetrachloro-3,6-diphenylglycoluril. *Biochemistry* 17:4807–4817
- Nichols EJ, Kannagi R, Hakomori SI, Krantz MJ, Fuks A (1985) Carbohydrate determinants associated with carcinoembryonic antigen (CEA). *J Immunol* 135:1911–1913
- Oikawa S, Nakazato H, Kosaki G (1987a) Primary structure of human carcinoembryonic antigen (CEA) deduced from cDNA sequence. *Biochem biophys Res Commun* 142:511–518
- Oikawa S, Imajo S, Noguchi T, Kosaki G, Nakazato H (1987b) The carcinoembryonic antigen (CEA) contains multiple immunoglobulin-like domains. *Biochem biophys Res Commun* 144:634–642
- Rogers GI (1983) Carcinoembryonic antigens and related glycoproteins. Molecular aspects and specificity. *Biochim Biophys Acta* 695:227–249
- Scatchard G (1949) The attractions of proteins for small molecules and ions. *Ann NY Acad Sci* 51:660–672
- Steinsträsser A, Schwarz A (1986) Quality control of iodinated monoclonal antibodies. *Nucl Med* 25:A75–76
- Steinsträsser A, Schwarz A, Kuhlmann L (1987) Monoclonal antibodies: Optimising of labelling procedures by quality control methods. Third European Symposium on Radiopharmacy and Radiopharmaceuticals; Kristensen K, Dige-Peterson H, Marquersen J (eds) *Elsinore, Abstract*, pp 68–69, Elsinore, Denmark, Isotope pharmacy bronshoj, May 1–4, 1987
- Towbin T, Gordon J (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications. *Proc Nat Acad Sci USA* 76:4350–4354
- Walker WHC (1977) The scatchard plot in immunometric assay. *Clin Chem* 23:588–590

Received March 4, 1988 / June 20, 1988

## A brief its use

J.A.J. Carrel  
Department  
P.O. Box 96  
Division of  
Department

**Abstract.**  
that can  
gas in the  
is describe  
6 views of  
50  $\mu\text{Sv}$  wa  
compariso  
**Key words**  
equivalent

During the  
come wide  
it is used  
nary emb  
tively. For  
and sensit  
the tradit  
lation stud  
1985; Free  
The re  
is presume  
is et al. 19  
ICRP 52  
to 800  $\mu\text{G}$   
rived from  
of counts  
to another  
gamma ca  
would be  
for a spe  
a given  
the total  
radioactiv  
dose equiv  
late the tr  
administer  
Before  
number of  
half life  
gas is add  
to determ  
tient. In a  
the total

Offprint rec